

EFFECT OF THERANOSTIC PHOSPHOLIPIDIC SRCL2/SRI2AND β -D-FRUCTOPYRANOSE BASED NANOPARTICLES ON CELL VIABILITY AND UPTAKE

Valentina Boscaro¹, Domenica Marabello², Paola Antoniotti², Paola Benzi², Alessandro Barge¹, Fabio Beccari², Carlo Canepa², Elena Peira¹, Margherita Gallicchio¹

¹Dipartimento di Scienza e Tecnologia del Farmaco - Università degli Studi di Turin, Turin - Italy, ²Dipartimento di Chimica - Università degli Studi di Turin, Turin - Italy

Introduction: Nanoparticles can be engineered with precise sizes, shapes, composition and with surface targeting ligands that allow a specific interaction with target cells. The aim of this work was to analyze the effect of non-toxic and low-cost Sr/I-based nanoparticles, in order to develop a new ⁸⁹Sr- and/or ¹³¹I-based radiotherapy method on cell lines. In particular we analyzed the effect on cellular uptake and cell viability of compound 1 (⁸⁹Sr(fructose)₂Cl₂·H₂O) and compound 2 (⁸⁹Sr(fructose)₂I₂ and Sr(fructose)₂¹³¹I₂), on a non-tumour cell line, hTERT-HME1, and a tumour cell line, HT-29. The compounds studied exhibit some features that could be interesting for biomedicine applications, such as the biocompatibility due to their non-toxic components and the significant SH emission, that can permit exploitation for in vitro bio-imaging.

Materials and methods: We tested the effect of nanoparticles encapsulated in a hydrophilic shell of mPEG-DSPE that increases their biocompatibility on non-tumour and tumour cell lines, hTERT-HME1 and HT-29, respectively. In particular, we analyzed the cellular uptake of nanoparticles and the effect on cell viability. To evaluate the effect of nanoparticles on cell viability, we treated hTERT-HME1 and HT-29 cell lines with five different nanoparticles concentrations, and cell viability was assessed using ATP content and the CellTiter-Glo® Luminescent Cell Viability Assay (Promega).

Results: hTERT-HME1 cell line showed sensitivity to compound 1 already at the lowest concentration, while compound 2 reduced hTERT-HME1 cell viability in a dose-dependent manner; HT-29 viability was reduced of about 70% and 50% only at the highest concentration of nanoparticles of compounds 1 and 2, respectively. The nanoparticles covered with mPEG-DSPE can be detected for few days in a biological medium and can be internalized into hTERT-HME1 and HT-29 cells.

Discussion and conclusions: In this work we have analyzed the effect of Sr/I-based nanoparticles, on cell uptake and viability. The nanoparticles formulation could be useful to concentrate the effect of ⁸⁹Sr and ¹³¹I radio-API in a specific tissue, allowing a selective activity of the activated compounds. The nanoparticles are biocompatible and are internalized into the cell; the viability of cell lines utilized is influenced by the presence of the nanoparticles themselves rather than their chemical composition.